
THE GEORGE
WASHINGTON
UNIVERSITY

WASHINGTON, DC

MP15-07: Timing of Ureteral Stent Removal After Ureteroscopy on Stent-Related Symptoms - A Validated Questionnaire Comparison of 3 and 7 Days Stent Duration


Heidenberg DJ¹, Helbig M¹, Grant C², Shoen E¹, Nauheim J³, Amdur R¹, Mufarrij P.¹

1. Department of Urology, George Washington University Hospital, Washington, DC

2. Department of Pediatric Urology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH

3. Department of Urology, Montefiore Medical Center, Bronx, NY


Study Design



Background	Patient Characteristics	n, (n%)
	Total Patients	n = 102
Age	Range	23-74
Gender		
	Male	51 (50)
	Female	51 (50)
Ethnicity		
	Caucasian	65 (64)
	African American	28 (27)
	Asian American	3 (3)
	Hispanic	4 (4)
Stent Duration		
	3 days	61 (60)
	7 days	41 (40)
Previous Stone Surgery		
	Yes	36 (35)
	No	66 (65)
Stent Size		
	6x22	8 (8)
	6x24	54 (53)
	6x26	37 (36)
	6x28	3 (3)

Results

Comparing Stent Related Symptoms for 3 days vs 7 days



■ Group 1 (In Situ)

■ Group 1 (Removed)

■ Group 2 (In Situ)

■ Group 2 (Removed)